

A NEW OXIDATIVE DERIVATIZATION METHOD FOR THE INDIRECT SPECTRO-FLUORIMETRIC DETERMINATION OF PROCHLORPERAZINE MALEATE IN PHARMACEUTICAL PREPARATIONS

M.YE. BLAZHEYEVSKIY, YU.V. SKRYPYNETS, A.V. YEGOROVA, V.P. ANTONOVICH

References:

1. Maryadele O.J. 14th ed. Whitehouse Station, New Jersey: Merck and Co., Inc; 2006. The Merck Index: An Encyclopedia of chemicals, drugs and biologicals, p. 1060.
2. U.S. Pharmacopoeia. NF Vol. II. The United states pharmacopeial convention, Twinbrook pathway Rockvillae, 2013. 2635 p.
3. British Pharmacopoeia. Vol. I. Stationary office. London Medicines and Healthcare product regulatory agency, 2010, 254-255.
4. European Pharmacopoeia 9th Edition – European Directorate for the Quality of Medicines (EDQM) Council of Europe, 67075 Strasbourg Cedex, France 2016. 4016 p.
5. Basavaiah K., Manjunatha J.M, Krishnamurthy G., Quantitation of pharmaceutically important phenothiazines by oxidimetry. *Farmaco*. 2000, 55 87-92. DOI: 10.1016/S0014-827X(99)00122-6
6. Kowalczyk-Marzec A., Kurzawa M., Szydłowska-Czerniak A., Szlyk E. Conductometric determination of phenothiazine derivatives by precipitation titration. *Chemia analityczna (Warsaw)*. 2002, 47(4), 613-618.
7. Basavaiah K., Swamy J.M. Titrimetric microdetermination of some phenothiazine neuroleptics with potassium hexacyanoferrate. *Talanta*. 1998, 47(1) 59-66. doi.org/10.1016/S0039-9140(98)00057-5
8. Shirsand S.B., Suresh S, Swamy P.V., Para M.S., Nagendra D.K. Formulation design of fast disintegrating tablets using disintegrant blends. *Indian J Pharm Sci*. 2010 Jan-Feb; 72(1), 130–133. DOI: 10.4103/0250-474X.62244
9. Shirsand S.B., Suresh S., Swamy P.V. Formulation design of novel fast disintegrating tablets of prochlorperazine maleate. *Int. J. Pharm. Tech. Res*. 2010, 2(1), 125-129.
10. Zare F., Mohammadloo Z.B. Survey Spectrophotometric Method for Determination of Phenothiazine Drug. *Nat Sci* 2016;14(10):1-5. doi:10.7537/marsnsj141016.01.
11. Upadhyay K., Asthana A., Tamrakar R.K. Sensitive spectrophotometric method for determination of some phenothiazine drugs. *Res. Chem. Intermediates*. 2015, 41(10), 7481-7495. DOI: 10.1007/s11164-014-1838-8.
12. Bhagwat G.B., Sonawane P.T., Bopinwar V.V. Determination of prochlorperazine maleate and pyridoxine hydrochloride in combined tablet dosage form by using first order derivative spectrophotometric method. *J. Sci. Tech*. 2012, 4(6). 262-266.
13. Bhagwat G.B., Wate S.P., Mundhey A.S. Development and Validation of UV Spectrophotometric method for estimation of prochlorperazine maleate and pyridoxine hydrochloride in tablet dosage form by UV using multi-component mode of analysis. *Indian drugs*. 2013, 50(3), 20-25.
14. Puzanowska-Tarasiewicz H., Karpin'ska J., and Kuzmicka L. Analytical Applications of Reactions of Iron(III) and Hexacyanoferrate(III) with 2,10-Disubstituted Phenothiazines. *Int. J. Anal. Chem*. 2009, Article ID 302696, 8 pages doi:10.1155/2009/302696
15. Theia'a N., Al-Talib S.M. Spectrophotometric determination of some phenothiazines using N-chlorosuccinimide. *Rafidain journal of science*, 2009, 20(4E), 27-37.
16. Revanasiddappa H.D., Veena, M.A. Spectrophotometric determination of some phenothiazine drugs using chloramine-T-iodine-starch. *J. Anal. Chem*. 2008, 63(2), 140-144. doi.org/10.1134/S1061934808020068
17. Basavaiah K. Indirect spectrophotometric determination of some biologically important phenothiazines using potassium dichromate, iron(II) and 1,10-phenanthroline. *Indian J. Chem. Tech*. 2004, 11(5):632-638
18. Aman T., Anwar J., Ahmad A., Latif L. Determination of five phenothiazines in pure and pharmaceutical preparations using vanadium pentoxide as a chromogenic reagent, *Anal. Lett*. 2003, 36(14), 2961-2974. DOI: 10.1081/AL-120026414
19. Basavaiah K., Swamy J. M. Application of potassium dichromate and iron-thiocyanate in the spectrophotometric investigations of phenothiazines. *Farmaco*. 2001, 56(8), 579–585. doi.org/10.1016/S0014-827X(01)01096-5
20. Aman T., Majeed Z., Kazi A.A., Khan I.U. Spectrophotometric Determination of Prochlorperazine Maleate in Pure and Pharmaceutical Preparations. *Mikrochim. Acta*. 2002, 138(1-2), 13-17. DOI: 10.1007/s006040200001
21. Gaiduk O.V., Pantaler R.P., Blank A.B. A New Catalytic Reaction for Determining Phenothiazine Derivatives. *J. Anal. Chem*. 2004, 59(7). 684-687. DOI: 10.1023/B:JANC.0000035284.55280.0b
22. Progress in Chemical Toxicology Vol. 5, Ed. Abraham Stolman. Elsevier, 2013, 406 p.

23. Nesměrák K., Červený V., Hraníček J., Rychlovský P. A spectrofluorimetric determination of phenothiazine derivatives after their photooxidation or chemical or electrochemical oxidation in a flow injection arrangement. *Microchem. J.* 2013, 106, 226–232. DOI: 10.1016/j.microc.2012.07.006
24. Yung G.J., Qu X.L., Ming, M., Qu Q.S., Wang C.Y., Zhu A.P., Hu A.Y. Trace measurement of phenothiazine drugs in tablets by micellar enhanced fluorophotometric method. *J. Fluoresc.* 2007, 17, 119–126. DOI: 10.1007/s10895-007-0165-5
25. Mohamed A.M., Abdelmageed O.H., Salem H., Nagy D.M., Omar M.A. Spectrofluorimetric determination of certain biologically active phenothiazines in commercial dosage forms and human plasma. *Luminescence.* 2013. 28(3):345-54. DOI: 10.1002/bio.2388.
26. Blazheyevskiy M.Ye. Application of derivatization by means of peroxy acid oxidation reaction in pharmaceutical analysis. *Farmacom.* 2016, 3, 28-40.
27. Yang Y., Peng Y., Zhao F., Zeng, B. Voltammetric determination of prochlorperazine and ethopropazine using a gold electrode modified with decanethiol SAM. *Sensors*, 2003. 3(11), 524-533. DOI: 10.3390/s31100524
28. Mielech-Łukasiewicz K., Puzanowska-Tarasiewicz H., Panuszko A. Electrochemical oxidation of phenothiazine derivatives at glassy carbon electrodes and their differential pulse and square-wave voltammetric determination in pharmaceuticals. *Anal. Lett.* 2008, 41(5), 789-805. DOI: 10.1080/00032710801935038
29. Kishorkumar L Mule. Rapid analytical method for assay determination for prochlorperazine edisylate drug substances by ultraperformance liquid chromatography. *Int. J. Curr. Pharm. Res.* 2017, 9(4), 118-122. DOI: 10.22159/ijcpr.2017v9i4.20973
30. Trawiński J., Skibiński R. Studies on photodegradation process of psychotropic drugs: a review. *Environ. Sci. Pollut. Res. Int.* 2017, 24(2), 1152-1199. DOI: 10.1007/s11356-016-7727-5
31. Tu Y.Y.; Hsieh M.M.; Chang S.Y. Sensitive detection of piperazinyl phenothiazine drugs by field-amplified sample stacking in capillary electrophoresis with dispersive liquid-liquid microextraction. *Electrophoresis*, 2015, 36 (21-22): 2828-2836. DOI: 10.1002/elps.201500260.
32. Dhabab J.M., Al-Ameri S.A., Taufeeq A.H. Separation and determination of trifluoperazine and prochlorperazine in pharmaceutical preparations by HPLC. *Arab Uni. for basic & app. sci.*, 2013, 13(1), 14-18.
33. Bhosale D., Nikalje A., Development and validation of RP-chromatography for simultaneous determination of paracetamol, caffeine, ergotamine and prochlorperazine in tablets. *Asian J. Pharm. Sci.*, 2013, 7(5): 364-370.
34. Tashiro M, Naito T, Kagawa Y, Kawakami J. Simultaneous determination of Prochlorperazine and its metabolites in human plasma using isocratic liquid chromatography tandem mass spectrometry. *Biomed Chrom.* 2012, 26(6),754-60. DOI: 10.1002/bmc.1725.
35. Yan M, Zhu Y.G., Li H.D., Chen B.M., Ma N., Lei Y.Q., Liu Y.P. Quantification of prochlorperazine maleate in human plasma by liquid chromatography-mass spectrometry: Application to a bioequivalence study. *J. Chrom. B. Analyt. Technol. Biomed Life Sci.* 2009, 877: 3243-3247.
36. Khanage S.G., Mohite P.B., Dudhade M.R. and Deshmukh V.K. Separation and Determination of Prochlorperazine Maleate and Pyridoxine Hydrochloride in Tablet Formulation by RP-HPLC. *Austin J Anal Pharm Chem.* 2014, 1(3): 1012.
37. Venkatesh D., Sundaram S., Meyyanathan S., Muralidharan S., Shanmugam R., Kannan E., Bhojraj S., Santhi S. Bioavailability Studies on Developed Prochlorperazine Maleate Sustained Release Tablets by HPLC, *JBABM*, 2009, 1: 54-57. DOI:10.4172/1948-593X.1000012
38. Patel Vishvas D., Patel Paresh U. Development and validation of reverse phase high performance liquid chromatography method for simultaneous estimation of betahistine dihydrochloride and prochlorperazine maleate in tablet dosage form. *Int. J. Pharm. Drug. Anal.* 2014, 2(4). 369-374.
39. Angela Lew, Qiufeng Gao & Lori H. Takahashi (2011) Analysis and identification of prochlorperazine impurities and forced degradation products using a stability-indicating hplc method, *Journal of Liquid Chromatography & Related Technologies*, 34:8, 634-651, DOI: 10.1080/10826076.2011.558650
40. Heiko Hayen and Uwe Kars Analysis of phenothiazine and its derivatives using LC/Electrochemistry/MS and LC/Electrochemistry/Fluorescence. *Anal. Chem.* 2003, 75(18), 4833-4840. DOI: 10.1021/ac0346050
41. Blazheyevskiy M.Ye. Application of derivatization by means of peroxy acid oxidation and perhydrolysis reactions in pharmaceutical analysis. Lviv: Ivan Franko National University of Lviv, 2017. 106 p.